



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2003

2,2,4,4-Tetramethylcyclobutane-1-one-3-thione

Heimgartner, Heinz ; Mlostoń, Grzegorz

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-80023>


Book Section

Published Version

Originally published at:

Heimgartner, Heinz; Mlostoń, Grzegorz (2003). 2,2,4,4-Tetramethylcyclobutane-1-one-3-thione. In: Paquette, Leo A; Rigby, James H; Crich, David; Wipf, Peter. Electronic Encyclopedia of Reagents in Organic Synthesis (e-EROS). Chichester: J. Wiley Sons, 1-14.

2,2,4,4-Tetramethylcyclobutane-1-one-3-thione



[10181-59-6]

C₈H₁₂OS

(MW 156.25)

(reagent used as a stable thioketone and a model for the comparison of the reactivities of strained carbonyl and thiocarbonyl groups)

Alternate Name: 2,2,4,4-tetramethyl-3-thioxocyclobutanone, 2,2,4,4-tetramethyl-3-thio-1,3-cyclobutanedione.

Physical Data: mp 57–58 °C.

Solubility: soluble in chlorinated hydrocarbons, hydrocarbons, ethers, alcohols, pyridine, and other commonly used organic solvents; insoluble in water.

Form Supplied in: red crystals, commercially not available.


Analysis of Reagent Purity: ¹H-NMR, ¹³C-NMR, single crystal X-ray analysis.¹

Preparative Methods: the title reagent can be prepared by thionation of 2,2,4,4-tetramethylcyclobutane-1,3-dione with P₄S₁₀ in pyridine at 110 °C.^{2–4}


Purification: column chromatography (silica gel, hexane/dichloromethane); sublimation.

Handling, Storage, and Precautions: camphoraceous odor; store in a refrigerator, safe in handling.

Oxidation Reaction. Treatment of 2,2,4,4-tetramethyl-3-thioxocyclobutanone (**1**) with *m*-CPBA in diethyl ether (or dichloromethane) at 0 °C gives the colorless sulfine (thioketone *S*-oxide, **2**) in 80% yield (eq 1).⁵ The same product is reported to be formed by sensitized photochemical oxidation of **1** with O₂/methylene blue in chloroform.⁶




The chlorination of **1** with gaseous chlorine in tetrachloromethane at 0 °C⁷ or with phosphorus pentachloride in boiling tetrachloromethane⁸ affords the stable α -chloro sulfanyl chloride (**3**) (eq 2). Phosphorus pentachloride can be replaced by sulfuryl chloride (SO₂Cl₂).




Sulfur dichloride in tetrachloromethane adds easily to **1** to give disulfanyl chloride **4** along with trisulfane **5** as a side product.^{8,9}


In an analogous reaction, disulfur dichloride and **1** react slowly and tetrasulfane **6** is formed as the exclusive product (eq 3).⁸




Reactions with Organometallics. Treatment of a mixture of **1** and alkyl bromides in THF with magnesium leads chemoselectively to 3-alkylthio-2,2,4,4-tetramethylcyclobutanones in a thiophilic addition of the in situ generated Grignard reagent.¹⁰ The thiophilic addition is also observed with methyllithium, and quenching with TMSCl results in the formation of bicyclo[1.1.0]butane (**7**) (eq 4).¹¹




Reactions with Carbenes and Carbenoids. Photochemical decomposition of diazomethane in the presence of **1** in an argon matrix at 10 K yields thiocarbonyl *S*-methylide (**8**) via addition of methylene to the sulfur atom.¹² While increasing the temperature, **8** undergoes a 1,3-dipolar electrocyclicization to give thiirane **9** (eq 5).




Under PTC conditions, involving trichloromethane and aqueous sodium hydroxide, **1** is converted into the *gem*-dichlorothiirane (**10**), which subsequently undergoes desulfurization and, finally, the C=O group is transformed into a stable *gem*-dichlorooxirane (**11**) (eq 6).¹³



When Seyferth's reagent [(phenyl)(trichloromethyl)mercury] is decomposed in boiling benzene in the presence of **1**, 84% of thiirane **10** is obtained. Through the use of (phenyl)(trifluoromethyl)mercury and sodium iodide under the same conditions, **1** is converted in 50% yield into the difluoro analog of **10**.¹⁴ Unexpectedly, difluorocarbene generated from bis(trifluoromethyl)cadmium in trichloromethane at $-20 \rightarrow 20^\circ\text{C}$ reacts with 2 equiv of **1** to give, in addition to the difluorothiirane derivative, 2,2-difluoro-1,3-dithiolane (a so-called Schönberg product, cf. ref 15) as the product of the interception of the intermediate thiocarbonyl *S*-difluoromethylide by **1**.¹⁶ In contrast to halogenated carbenes, the nucleophilic dimethoxycarbene, generated by thermolysis of **12**, reacts with **1** to give the ring enlarged products **13** and **14** in favor of **13** (eq 7).¹⁷ Remarkably, no dimethoxy thiirane derivative of **1** can be detected.




The decomposition of dimethyl diazomalonate with $\text{Rh}_2(\text{OAc})_4$ in toluene at 50°C in the presence of **1** leads to two products **15** and **16**, which are formed via an intermediate thiocarbonyl ylide (eq 8). Whereas 1,3-dipolar electrocyclization affords a thiirane which spontaneously extrudes sulfur to give **15**, the competitive 1,5-dipolar electrocyclization yields 1,3-oxathiole **16**.¹⁸




Reactions with 1,3-Dipoles

Diazo Compounds


Diazomethane reacts immediately with **1** dissolved in diethyl ether in a regioselective manner to give 2,5-dihydro-1,3,4-thiadiazole (**17**) as a relatively stable solid.^{19,20} This compound is a superior precursor of the reactive thiocarbonyl *S*-methylide (**8**, eq 9) as it smoothly eliminates nitrogen at 45 °C. In the absence of an appropriate interceptor, thiirane **9** is formed.²¹ In the presence of electron-deficient dipolarophiles, diverse five-membered spiro-heterocycles (**18**) are formed via 1,3-dipolar cycloaddition (eq 9).^{22,23} Representative examples of dipolarophiles leading to **18** in high yield are DMAD, *N*-phenyl maleinimide, chloral, dimethyl azodicarboxylate, adamantanethione, as well as **1**.²¹




The formation of 1,3-oxathiolane with chloral and 1,3-dithiolanes with cycloaliphatic thioketones occurs regioselectively to yield the sterically less hindered products.¹⁵ On the other hand, aromatic thioketones, e.g., thiobenzophenone or 9*H*-fluorene-9-thione, intercept **8** to give comparable amounts of both regioisomeric adducts. Usually, stereoisomeric dipolarophiles such as fumaronitrile and maleonitrile as well as dimethyl fumarate and maleate form **18** in a stereoselective manner. However, in the case of extremely electron-poor dipolarophiles, e.g., dimethyl 1,2-dicyanofumarate or (*E*)-1,2-bis(trifluoromethyl)ethylene-1,2-dicarbonitrile, non-stereospecific formations of the corresponding tetrahydrothiophenes are described.^{24,25} This result is interpreted in terms of a stepwise reaction mechanism with a zwitterion as the key intermediate.^{25,26} Alternatively, this intermediate can cyclize to form seven-membered ketenimines of type **20**. With R = CF₃, this product can be isolated in a crystalline form, whereas in the case of R = CN, stable lactam **21** is obtained only after addition of water (eq 10).



2-Diazopropane reacts with **1** at 0 °C to yield the stable 2,2-dimethyl analog of **17**, which is used as a precursor of the *S*-isopropylidene of **1**.²⁷ The 2,2-diphenyl analog of **17**, prepared from **1** and diphenyldiazomethane, extrudes nitrogen spontaneously leading to the corresponding thiirane.²⁷ Similarly, EDA and **1** react at rt to give a thiirane-2-carboxylate, which can be desulfurized to yield an ethyl cyclobutylidene acetate.²⁸ The reactions of **1** with less reactive diazocompounds such as diazoketones, diazoamides, and diazoesters are accelerated by addition of LiClO₄, and thiiranes and/or 1,3-oxathioles are obtained.^{28–31} Products of both types are formed in the case of α -diazoamide (**22**) (eq 11).




With a more complex diazo compound containing amino and ester groups, products resulting from 1,3- and 1,5-dipolar electrocyclization are obtained.³² ‘Two-fold extrusion’ reactions with **1** and diverse diazo compounds are reported to be aimed at the preparation of sterically crowded ethylenes.^{33,34} An example with bis(*tert*-butyl)diazomethane (**25**) is shown in eq 12.




Organic Azides

The 1,3-dipolar cycloadditions of **1** with organic azides require elevated temperatures. Typical products of the reactions of thioketones are *N*-substituted imines, which are formed after decomposition of the initially formed 1,2,3,4-thiazotriazole derivatives.^{35,36} However, in the case of **1** and aryl azides, the intermediate thiocarbonyl *S*-imides (**28**) intercept **1** to give dispiro-1,4,2-dithiazoles (**29**) (eq 13).³⁷




The analogous reaction of **1** with benzyl azide leads to a rearrangement of the corresponding *N*-benzyl derivative of **28** via a 1,4-H shift.³⁸ In some cases, e.g., methyl azidoacetate, the sulfur atom extruded in the course of the reaction is intercepted by **1** to give a thiocarbonyl *S*-sulfide (thiosulfine, **33**), which subsequently undergoes a 1,3-dipolar cycloaddition with **1** yielding 1,2,4-trithiolane (**34**).³⁶ Thiaziridine (**31**) formed by ring-closure of **30** is a plausible sulfur donor (eq 14).



In a three-component reaction with **1** and fumaronitrile dissolved in phenyl azide, the *N*-phenyl substituted thiocarbonyl *S*-imide (**28**, Ar = Ph) cycloadds to fumaronitrile to give the corresponding 1,2-thiazolidine derivative.³²


Nitrones

1,3-Dipolar cycloadditions of nitrones with **1** occur regioselectively and 1,4,2-oxathiazolidines of type **36** are obtained in high yield. Aldonitrones are more reactive than ketonitrones.³ When stored in CDCl₃ solution at 25 °C, cycloreversion takes place and, in the case of **35**, the equilibrated system contains 4% of **1** with R = H and 14% with R = Ph, respectively (eq 15).^{39,40}



Azomethine Ylides


Various azomethine ylides, generated by different methods, can be trapped by **1** to give spirocyclic 1,3-thiazolidines. Thermolysis of *cis*-aziridines of type **37** in the presence of **1** occurs stereoselectively to give *trans*-configured cycloadducts (**38**) (eq 16).⁴¹ Reactions carried out with *trans*-aziridines lead to *cis*-substituted 1,3-thiazolidines.



Using the desilylation methodology for the generation of azomethine ylides, 2,4-unsubstituted *N*-benzyl as well as *N*-unsubstituted 1,3-thiazolidines of type **38** are prepared.^{42,43}

Thiocarbonyl Ylides


Thiocarbonyl ylides generated by thermolysis of 2,5-dihydro-1,3,4-thiadiazoles can be intercepted by **1** to give 1,3-dithiolane derivatives (so-called Schönberg products).⁴⁴ Both aromatic and cycloaliphatic *S*-methylides undergo [2+3]-cycloadditions with **1** regioselectively. Whereas **39** yields the sterically crowded adduct **40**,⁴⁵ adamantane *S*-methylide (**41**) affords the less hindered 1,3-dithiolane (**42**) (eq 17).⁴⁶ However, the reaction of **8**, the *S*-methylide of **1**, with thiobenzophenone results in a mixture of **40** and the opposite regioisomer.⁴⁵




Thiocarbonyl *S*-imides

In eq 13, the interception of a thiocarbonyl *S*-imide generated in situ from **1** and organic azides is outlined.³⁷ Similarly, thiocarbonyl *S*-imide (**30**, eq 14) undergoes a 1,3-dipolar cycloaddition with **1** in competition with the formation of the unstable thiaziridine **31**, and the corresponding 2,5-dispiro-1,4,2-dithiazolidine (**43**) and imine **32** are obtained (eq 18).³⁶ A similar reaction with methyl dithiobenzoate, in which along with 1,2,4-trithiolane (**34**)

a 1,4,2-dithiazolidine of type **43** is formed, is also reported.⁴⁷




The stable, sterically crowded hexafluorothioacetone *S*-imide (**44**) reacts smoothly with aromatic thioketones, but the reaction with **1** requires elevated temperatures.^{48,49} Unexpectedly, the product isolated after 3 h at 100 °C is 1,4,2-dithiazolidine (**47**). Its formation can be rationalized by the isomerization of **44** to give thiaziridine **45** being in an equilibrium with thionitrone **46**, which then traps **1** to yield **47** (eq 19).




Thiocarbonyl *S*-oxides (Sulfines)

Cycloaliphatic sulfines such as **2** or **48** react with **1** in chloroform at 80 °C to give the stable 1,2,4-oxadithiolanes of type **49** in a regioselective [2+3] cycloaddition (eq 20).⁵⁰




In contrast, the analogous cycloadducts of aromatic sulfines and **1** cannot be isolated. Instead, 1,2,4-trithiolane (**51**) is obtained in the reaction of thiobenzophenone *S*-oxide (**50**) with 2 equiv of **1** (eq 21).⁵¹ In this multi-step reaction, a sulfur-transfer leading to a thiocarbonyl *S*-sulfide is proposed as a key step. The formation of equimolar amounts of 2,2,4,4-tetramethylcyclobutane-1,3-dione

(**52**) indicates that in this system **1** acts as a sulfur donor. In a three-component system containing **50**, **1**, and (*E*)-cyclooctene, an episulfidation of the strained alkene is observed.⁵²




Thiocarbonyl *S*-sulfides (Thiosulfines)


The formation of the symmetrical 1,2,4-trithiolane (**34**, eq 14) is the evidence for the intermediacy of thiosulfine **33** formed by a sulfur transfer from thiaziridine **31** to **1**. The mixed 1,2,4-trithiolane (**51**) (eq 22) can result either from the [2+3]-cycloaddition of thiobenzophenone *S*-sulfide (**53**) with **1** or from **33** with thiobenzophenone. Both pathways are conceivable for the formation of **51** in a three-component reaction including **1**, thiobenzophenone, and phenyl azide.⁵³ Tetraphenyl-1,2,4-trithiolane (**54**) undergoes a [2+3]-cycloreversion and releases **53**, which is trapped by **1** to afford **51** (eq 22). The latter is stable under the reaction conditions.⁵¹




Isomerizations. The isomerization of **1** to thiolactone **55** occurs in DMSO solution in the presence of NaH.² It is reported that photolysis of **1** in methanol also affords **55**.⁵⁴ The reaction of **1** with (trifluoromethyl)trimethylsilane (Ruppert's reagent) in THF in the presence of fluoride yields the trifluoromethylated thietane **56** (eq 23).⁵⁵ A plausible mechanism of the transformation is a ring-opening/ring-closure process initiated by nucleophilic addition of trifluoromethanide to the carbonyl group of **1**.




Photolysis of **1** in benzene is reported to give another isomer with the structure **57** along with the ring-enlarged 3-oxodithiolactone (**58**) (eq 24).⁵⁶




Miscellaneous. The sulfur-rich heterocycles **59** and **60** are produced when **1** in acetone is treated with elemental sulfur (S_8) in the presence of catalytic amounts of sodium thiophenolate (eq 25).⁵⁷ The type of the product depends on the ratio of **1** and S_8 used in the reaction.




The related hexathiepane **62** can be prepared from the acetylated trisulfane **61**, obtained in two steps from **1**, by treatment with morpholine in THF solution (eq 26).⁸




Lewis acid-catalyzed reactions of **1** with oxiranes lead to 1,3-oxathiolanes and/or 1,3-dithiolanes. Whereas cyclohexene oxide (**63**) yields a mixture of **64** and **65**, only the corresponding 1,3-dithiolane is obtained in the case of cyclopentene oxide (eq 27).⁵⁸ The formation of 1,3-dithiolanes is evidenced to involve the corresponding thiirane formed in situ by cleavage of the 1,3-oxathiolane ring. The structures of the products show that the heterocycle is formed with inversion of the configuration of one oxirane C-atom. This explanation is supported by the results obtained with **1** and *cis*- and *trans*-2,3-dimethyloxirane, respectively.⁵⁹ In this case, dispirocyclic 1:2 adducts are also formed by involvement of the carbonyl group of **1**.



A straightforward conversion of 2-unsubstituted imidazole 3-oxides (**66**) into the corresponding imidazole-2-thiones (**67**) is achieved by the reaction with **1** in dichloromethane at rt (eq 28).⁶⁰ This sulfur-transfer reaction can be applied to other azole *N*-oxides with an unsubstituted carbon atom next to the *N*-oxide position.



The reaction of amino-9-fluorenylideneborane (**68**) with **1** occurs chemo- and regioselectively at the carbonyl group to give the 1,2-oxaboretane (**69**) (eq 29).⁶¹



The irradiation of **1** in 2-propanol in the presence of oxygen at wavelengths > 400 nm results in the desulfurization of **1** to yield the parent dione **52**. Most likely, sulfur monoxide is a side product.²

1. Shirrell, C. D.; Williams, D. E., *Acta Crystallogr., Sect. B* **1974**, *30*, 1974–1978.
2. Elam, E. U.; Davis, H. E., *J. Org. Chem.* **1967**, *32*, 1562–1564.
3. Worman, J. J.; Shen, M.; Nichols, P. C., *Can. J. Chem.* **1972**, *50*, 3923–3926.
4. Black, D. S. C.; Watson, K. G., *Aust. J. Chem.* **1973**, *26*, 2491–2504.
5. Zwanenburg, B.; Wagenaar, A.; Thijs, L.; Strating, J., *J. Chem. Soc., Perkin Trans. I* **1973**, 73–75.
6. Sundari, B.; Ramamurthy, V., *Indian J. Chem., Sect. B* **1984**, *23*, 498–501.
7. Koch, K. N.; Mloston, G.; Senning, A., *Eur. J. Org. Chem.* **1999**, 83–86.
8. Mloston, G.; Majchrzak, A.; Senning, A.; Sjøtofte, I., *J. Org. Chem.* **2002**, *67*, 5690–5695.
9. Linden, A.; Majchrzak, A.; Cavegn, J.; Mloston, G.; Heimgartner, H., *Acta Crystallogr., Sect. C* **2002**, *58*, 480–484.
10. Masson, J.; Metzner, P.; Vialle, J., *Tetrahedron* **1977**, *33*, 3089–3095.
11. Gassman, P. G.; Mullins, M. J., *Tetrahedron Lett.* **1980**, *21*, 2219–2222.
12. Mloston, G.; Romanski, J.; Schmidt, C.; Reisenauer, H. P.; Maier, G., *Chem. Ber.* **1994**, *127*, 2527–2530.
13. Mloston, G.; Romanski, J.; Swiatek, A.; Heimgartner, H., *Helv. Chim. Acta* **1999**, *82*, 946–956.
14. Mloston, G.; Romanski, J.; Heimgartner, H., *Heterocycles* **1999**, *50*, 403–410.
15. Huisgen, R.; Mloston, G.; Polborn, K.; Sustmann, R., *Chem. Eur. J.* **2003**, *9*, 2256–2263.
16. Haiges, R., Ph.D. thesis, University of Wuppertal, Germany, 2000; http://pub.bib.uni-wuppertal.de/edoos/fachbereiche/09_chemie
17. Dawid, M.; Mloston, G.; Warkentin, J., *Chem. Eur. J.* **2002**, *8*, 2184–2187.
18. Mloston, G.; Romanski, J.; Heimgartner, H., *Pol. J. Chem.* **2002**, *76*, 551–555.
19. Diebert, C. E., *J. Org. Chem.* **1970**, *35*, 1501–1505.
20. Huisgen, R.; Penelle, J.; Mloston, G.; Buyle Padias, A.; Hall, H. K., *J. Am. Chem. Soc.* **1992**, *114*, 266–274.
21. Huisgen, R.; Mloston, G.; Fulka, C., *Heterocycles* **1985**, *23*, 2207–2217.
22. Mloston, G.; Heimgartner, H., *Pol. J. Chem.* **2000**, *74*, 1503–1532.
23. Mloston, G.; Heimgartner, H., In *The Chemistry of Heterocyclic Compounds, Vol. 59: Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products*; Padwa, A.; Pearson, W. H., Eds.; Wiley: New York, 2002, pp 315–360.
24. Huisgen, R.; Mloston, G.; Giera, H.; Langhals, E., *Tetrahedron* **2002**, *58*, 507–519.
25. Huisgen, R.; Mloston, G.; Langhals, E.; Oshima, T., *Helv. Chim. Acta* **2002**, *85*, 2668–2685.
26. Huisgen, R.; Mloston, G.; Langhals, E., *Helv. Chim. Acta* **2001**, *84*, 1805–1820.

27. Mloston, G.; Huisgen, R.; Giera, H., *Tetrahedron* **2002**, 58, 4185–4193.
28. Kägi, M.; Mloston, G.; Heimgartner, H., *Pol. J. Chem.* **1998**, 72, 678–687.
29. Kägi, M.; Linden, A.; Mloston, G.; Heimgartner, H., *Helv. Chim. Acta* **1996**, 79, 855–874.
30. Kägi, M.; Linden, A.; Mloston, G.; Heimgartner, H., *Helv. Chim. Acta* **1998**, 81, 285–302.
31. Kelmendi, B.; Mloston, G.; Heimgartner, H., *Heterocycles* **2000**, 52, 475–482.
32. Mloston, G.; Romanski, J.; Linden, A.; Heimgartner, H., *Helv. Chim. Acta* **1996**, 79, 1305–1314.
33. Garratt, P. J.; Payne, D.; Tocher, D. A., *J. Org. Chem.* **1990**, 55, 1909–1915.
34. Freund, W.; Huenig, S., *J. Org. Chem.* **1987**, 52, 2154–2161.
35. Guziec, F. S.; Moustakis, C., *J. Chem. Soc., Chem. Commun.* **1984**, 63–64.
36. Mloston, G.; Romanski, J.; Linden, A.; Heimgartner, H., *Pol. J. Chem.* **1996**, 70, 880–890.
37. Mloston, G.; Romanski, J.; Linden, A.; Heimgartner, H., *Helv. Chim. Acta* **1993**, 76, 2147–2154.
38. Mloston, G.; Romanski, J.; Linden, A.; Heimgartner, H., *Helv. Chim. Acta* **1995**, 78, 1067–1078.
39. Huisgen, R.; Fisera, L.; Giera, H.; Sustmann, R., *J. Am. Chem. Soc.* **1995**, 117, 9671–9678.
40. Giera, H.; Huisgen, R., *Liebigs Ann./Recueil* **1997**, 1685–1689.
41. Mloston, G.; Urbaniak, K.; Linden, A.; Heimgartner, H., *Helv. Chim. Acta* **2002**, 85, 2644–2656.
42. Gebert, A.; Linden, A.; Mloston, G.; Heimgartner, H., *Pol. J. Chem.* **2003**, 77, 157–167.
43. Gebert, A.; Linden, A.; Mloston, G.; Heimgartner, H., *Pol. J. Chem.* **2003**, 77, 867–879.
44. Mloston, G.; Huisgen, R., *Heterocycles* **1985**, 23, 2201–2206.
45. Huisgen, R.; Li, X.; Mloston, G.; Fulka, C., *Eur. J. Org. Chem.* **2000**, 1695–1702.
46. Mloston, G.; Huisgen, R.; Polborn, K., *Tetrahedron* **1999**, 55, 11475–11494.
47. Mloston, G.; Romanski, J.; Linden, A.; Heimgartner, H., *Helv. Chim. Acta* **1995**, 78, 1499–1510.
48. Mloston, G.; Celeda, M.; Roesky, H. W.; Parisini, E.; Ahlemann, J. T., *Eur. J. Org. Chem.* **1998**, 459–465.
49. Mloston, G.; Lesniak, S.; Linden, A.; Roesky, H. W., *Tetrahedron* **2000**, 56, 4231–4238.
50. Huisgen, R.; Mloston, G.; Polborn, K.; Sustmann, R.; Sicking, W., *Liebigs Ann./Recueil* **1997**, 179–185.
51. Huisgen, R.; Mloston, G.; Polborn, K.; Palacios-Gambra, F., *Liebigs Ann./Recueil* **1997**, 187–192.
52. Adam, W.; Bargon, R. M.; Mloston, G., *Eur. J. Org. Chem.* **2003**, 4012–4015.
53. Mloston, G.; Heimgartner, H., *Helv. Chim. Acta* **1995**, 78, 1298–1310.
54. Muthuramu, K.; Ramamurthy, V., *Chem. Lett.* **1981**, 1261–1264.
55. Mloston, G.; Prakash, G. K. S.; Olah, G. A.; Heimgartner, H., *Helv. Chim. Acta* **2002**, 85, 1644–1658.
56. Rao, N. B.; Chandrasekhar, J.; Ramamurthy, V., *J. Org. Chem.* **1988**, 53, 745–751.
57. Huisgen, R.; Rapp, J.; Huber, H., *Liebigs Ann. Recueil* **1997**, 1517–1524.
58. Blagoev, M.; Linden, A.; Heimgartner, H., *Helv. Chim. Acta* **1999**, 82, 2316–2335.
59. Blagoev, M.; Linden, A.; Heimgartner, H., *Helv. Chim. Acta* **2000**, 83, 3163–3178.
60. Mloston, G.; Gendek, T.; Heimgartner, H., *Helv. Chim. Acta* **1998**, 81, 1585–1595.

61. Channareddy, S.; Glaser, B.; Mayer, E. P.; Nöth, H.; Helm, S. W., *Chem. Ber.* **1993**, *126*, 1119–1125.

Heinz Heimgartner
*Organisch-Chemisches Institut, Universität Zürich,
Switzerland*

Grzegorz Mloston
University of Lodz, Lodz, Poland